

























## Booster Dose

A single Janssen COVID-19 Vaccine booster dose (0.5 mL) may be administered at least 2 months after primary vaccination with the Janssen COVID-19 Vaccine, to individuals 18 years of age and older.

A single booster dose of the Janssen COVID-19 Vaccine (0.5 mL) may be administered to individuals 18 years of age and older as a heterologous booster dose following completion of primary vaccination with another authorized or approved COVID-19 vaccine. The dosing interval for the heterologous booster dose is the same as that authorized for a booster dose of the vaccine used for primary vaccination.

### **3 DOSAGE FORMS AND STRENGTHS**

Janssen COVID-19 Vaccine is a suspension for intramuscular injection. A single-dose is 0.5 mL.

### **4 CONTRAINDICATIONS**

#### **4.1 Severe Allergic Reactions**

Do not administer the Janssen COVID-19 Vaccine to individuals with a known history of severe allergic reaction (e.g., anaphylaxis) to any component of the Janssen COVID-19 Vaccine [*see Description (13)*].

#### **4.2 Thrombosis with Thrombocytopenia**

Do not administer the Janssen COVID-19 Vaccine to individuals with a history of thrombosis with thrombocytopenia following the Janssen COVID-19 Vaccine or any other adenovirus-vectored COVID-19 vaccine (e.g., AstraZeneca's COVID-19 vaccine which is not authorized or approved in the United States) [*see Warnings and Precautions (5.2)*].

### **5 WARNINGS AND PRECAUTIONS**

#### **5.1 Management of Acute Allergic Reactions**

Appropriate medical treatment used to manage immediate allergic reactions must be immediately available in the event an acute anaphylactic reaction occurs following administration of the Janssen COVID-19 Vaccine.

Monitor Janssen COVID-19 Vaccine recipients for the occurrence of immediate adverse reactions according to the Centers for Disease Control and Prevention guidelines (<https://www.cdc.gov/vaccines/covid-19/clinical-considerations/managing-anaphylaxis.html>).

#### **5.2 Thrombosis with Thrombocytopenia Syndrome (TTS)**

Reports to the Vaccine Adverse Events Reporting System (VAERS), a passive surveillance system, provide evidence for an increased risk of thrombosis with thrombocytopenia syndrome (TTS) with onset of symptoms approximately one to two weeks after administration of the Janssen COVID-19 Vaccine.
















































## 21 CONTACT INFORMATION

For general questions or to access the most recent Janssen COVID-19 Vaccine Fact Sheets, scan the QR code using your device, visit [www.janssencovid19vaccine.com](http://www.janssencovid19vaccine.com) or call the telephone numbers provided below.

QR Code	Fact Sheets Website	Telephone numbers
	<a href="http://www.janssencovid19vaccine.com">www.janssencovid19vaccine.com</a>	US Toll Free: 1-800-565-4008 US Toll: 1-908-455-9922

This Full EUA Prescribing Information may have been updated. For the most recent Full EUA Prescribing Information, please see [www.janssencovid19vaccine.com](http://www.janssencovid19vaccine.com).

Manufactured by:  
Janssen Biotech, Inc.  
a Janssen Pharmaceutical Company of Johnson & Johnson  
Horsham, PA 19044, USA



Revised: JAN/11/2022  
© 2021 Janssen Pharmaceutical Companies